RECOMMENDATIONS

Pullets

Starting period
week 1

		D1	D2	D3	D4	D5	D6	D7
SeaLyt	Water intake stimulation and rehydration	●	●					
Searup	Revive natural defences			●●	●●	●●		

Around the vaccination

		D-3	D-2	D-1	* vaccine day	D+1	D+2	D+3
Searup	Vaccines supplied by drinking water			●		●	●	
Searup	Vaccines supplied by injection or in the eyes			●		●	●	

* *According to the vaccination program*

Arrival of pullets
(including the first set up hours)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16
Time (in weeks)

Searup	Revive natural defences	● ● ● ● ● ● ● ●
		recommended dose: ***the day before vaccine and the 2 days after***
SeaLyt	In case of digestive troubles	*recommended dose:* ***during 3 days***
SeaLyt	In case of thermal stress	*recommended dose:* ***during 1 to 3 days***

● *single dose*
●● *double dose*

Local regulations should be consulted concerning the status of this product in the country of destination. Guide d'utilisation volaille - V1 du 03.12.15 EN

RECOMMENDATIONS

Laying hens

Standard laying curve of laying hens:



Transfer day
week 16

		D1	D2	D3	D4	D5	D6	D7
SeaLyt	Rehydration	●						
Searup	Revive natural defences		●●	●●	●●			

Egg production peak
around week 22

		D1	D2	D3	D4	D5	D6	D7
Searup	Help to maintain the production level	●	●	●				

Around the vaccination

		D-3	D-2	D-1	* vaccine day	D+1	D+2	D+3
Searup	Vaccines supplied by drinking water			●		●	●	
Searup	Vaccines supplied by injection or in the eyes			●		●	●	

* *According to the vaccination program*

Searup	Revive natural defences	● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ●
		recommended dose: ***once a week*** *+ in case of* ***immune depression*** *or* ***during risk periods***
DigestSea	Improve digestive functions	● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ●
		recommended dose: ***once a month*** *during* ***3 days:*** *0.3ml / liter / day*
SeaLyt	In case of digestive troubles or thermal stress	*recommended dose: during* ***1 to 3 days*** *+ in case of* ***risk period***

● *single dose*
●● *double dose*

Local regulations should be consulted concerning the status of this product in the country of destination. Guide d'utilisation volaille - V1 du 03.12.15 EN

Broilers

RECOMMENDATIONS



Local regulations should be consulted concerning the status of this product in the country of destination Guide utilisation volaille - V3 du 03-12-15 EN

RECOMMENDATIONS TO USE FOR POULTRY

***MSP**
Marine Sulfated Polysaccharides

SeaLyt

REDUCES DIGESTIVE TROUBLES
CONTAINS MSP* MUCIN FAVORING THE PROTECTION OF GUT MUCOSA

- Starter periods
- Heat stress
- Digestive troubles

Global application: single dose(•)**

Spark
1 spark 87g / 1000 liters
1 spark 45g / 500 liters
1 spark 16g / 200 liters

Searup

REINFORCES NATURAL DEFENSES
CONTAINS MSP* WITH IMMMUNOMODULATING PROPERTIES

- Supports vaccination programs
- Stress periods (starter periods, production peak, buildings changes...)

Global application: single dose(•)**

Spark
1 spark 80g / 1000 liters
1 spark 45g / 500 liters
1 spark 16g / 200 liters

Still
0.5 ml / liter
0.5 ml / 10 kg of animal live weight

DigestSea

IMPROVES DIGESTIVE FUNCTIONS, PROTECTS THE LIVER
CONTAINS MSP* WITH ANTI-HYPERLIPIDEMIC PROPERTIES

- Higher metabolic demand periods

Global application**

Still
0.3 to 1 ml / liter
0.3 to 1 ml / 10 kg of animal live weight

*** recommended daily doses*